

Relationship of Thyroid Hormone Levels to Levels of Polychlorinated Biphenyls, Lead, *p,p'*-DDE, and Other Toxicants in Akwesasne Mohawk Youth

Lawrence M. Schell,^{1,2} Mia V. Gallo,¹ Melinda Denham,¹ Julia Ravenscroft,¹ Anthony P. DeCaprio,³ and David O. Carpenter⁴

¹Department of Anthropology, University at Albany, State University of New York, Albany, New York, USA; ²Department of Epidemiology and Biostatistics, School of Public Health, University at Albany, State University of New York, Rensselaer, New York, USA; ³School of Public Health and Health Sciences, University of Massachusetts, Amherst, Massachusetts, USA; ⁴Institute for Health and the Environment, University at Albany, State University of New York, Rensselaer, New York, USA

BACKGROUND: It is well documented that acute exposure to high levels of persistent organic pollutants, such as polychlorinated biphenyls (PCBs), *p,p'*-dichlorophenyldichloroethylene (*p,p'*-DDE), and hexachlorobenzene (HCB), can affect human health including thyroid function. Chronic exposure to multiple toxicants is common but difficult to analyze, and most prior studies have focused on adults or newborns, creating a gap in our understanding of multitoxicant effects among adolescents.

OBJECTIVE: We investigated whether levels of PCBs, *p,p'*-DDE, HCB, mirex, lead, and mercury reflecting past chronic exposure are associated with alterations in levels of thyroid-stimulating hormone (TSH), triiodothyronine (T₃), total thyroxine (TT₄), and free thyroxine (FT₄) among older children and adolescents.

METHODS: The sample consists of youth from the Akwesasne Mohawk Nation (*n* = 232) who reside in proximity to several industries that have contaminated the local environment. We used multiple regression analysis to examine the effect of PCB groupings, *p,p'*-DDE, HCB, lead, and mercury on thyroid hormones after adjusting for sociodemographic covariates and controlling for all other toxicants.

RESULTS: Exposure to PCBs affects the thyroid hormone profile in adolescents. The group of persistent PCBs was positively associated with TSH but inversely related to FT₄. Nonpersistent PCBs were significantly and negatively related to FT₄ only. HCB was negatively associated with T₄, and lead was positively associated with T₃. Breast-fed adolescents had higher levels of persistent PCBs and *p,p'*-DDE but not of nonpersistent PCBs or any other toxicant when compared with non-breast-fed adolescents. Though having lower levels of persistent PCBs and *p,p'*-DDE, non-breast-fed adolescents exhibited significant relationships between persistent PCBs and TSH and FT₄, but breast-fed adolescents did not. It appears that PCBs from breast milk obscure the relationship between prenatal PCB exposure and thyroid function by adding random variation in PCB levels.

CONCLUSION: Our results demonstrate a reduction in thyroid function in adolescents in relation to their current serum levels of PCBs. These observations are consistent with the hypothesis that prenatal exposure to PCBs alters thyroid function in a long-lasting manner but does not exclude the possibility that postnatal exposure is influential also.

KEY WORDS: adolescents, hexachlorobenzene, mercury, mirex, Mohawk, Native American, PCBs, POPs, *p,p'*-dichlorophenyldichloroethylene, thyroid, thyroid hormones. *Environ Health Perspect* 116:806–813 (2008). doi:10.1289/ehp.10490 available via <http://dx.doi.org/> [Online 25 February 2008]

Exposure to persistent organic pollutants (POPs) such as polychlorinated biphenyls (PCBs), *p,p'*-dichlorophenyldichloroethylene (*p,p'*-DDE), a metabolite of *p,p'*-dichlorodiphenyltrichloroethane (*p,p'*-DDT), and hexachlorobenzene (HCB) is a global phenomenon. High levels of exposure to these and other potentially endocrine-disrupting toxicants affect human health, but the full extent of their impact remains an important area of study (Carpenter 2006; Daston et al. 2003; Kimbrough and Krouskas 2003; Koppe et al. 2006; Langer 2005). Experimental animal studies have demonstrated effects of POPs on endocrine system functioning, including thyroid function (Brucker-Davis 1998). These studies typically investigate large, single-toxicant exposures that are uncharacteristic of most human exposure patterns. Studies in humans have found relationships between exposure to specific POPs and lower thyroid

hormone levels as well as higher thyroid-stimulating hormone (TSH) levels (Koopman-Esseboom et al. 1994; Nagayama et al. 1998; Osius et al. 1999; Persky et al. 2001; Ribas-Fito et al. 2003; Rylander et al. 2006; Sauer et al. 1994; Wang et al. 2005; Zuurbier et al. 2006). Langer et al. (2005) have demonstrated that PCB exposure results in an increase in thyroid gland volume but an inverse relationship with TSH (2006). However, not all investigators have observed such associations, and as human studies have focused almost entirely on infants and adults (Meeker et al. 2006), the relationship of toxicants to thyroid hormone status in older children and adolescents is not apparent (Hagmar 2003).

Thyroid hormones are essential, regulating metabolism and promoting normal cardiovascular, reproductive, and nervous system functioning (Larsen et al. 2003). They are necessary for normal growth and brain development

(Porterfield and Hendrich 1993), and thus may represent a causal link between toxicants and observed effects on somatic growth and cognitive development.

The aim of the current investigation is to assess whether levels of PCBs indicative of a chronic exposure pattern are associated with alterations in levels of TSH, triiodothyronine (T₃), total thyroxine (TT₄), and free thyroxine (FT₄) among older children and adolescents. In addition to congener-specific PCB analyses, we also examined the effects of other common toxicants (*p,p'*-DDE, HCB, mirex, lead, and mercury) on thyroid hormone levels.

Methods

Setting. The study was conducted with mother–youth pairs who were members of the Akwesasne Mohawk Nation (Akwesasne), which spans the St. Lawrence River with territory in New York State and in Ontario and Quebec, Canada. Industrial development along the St. Lawrence River began in the 1950s and continues today, with several industrial complexes located near Akwesasne. A National Priority Superfund Site (General Motors Central Foundry Division) and two New York State Superfund Sites (Reynolds Metal Company and Aluminum Company of America) are located immediately upstream. In the 1990s, some local animal species were found to have levels of PCBs, *p,p'*-DDE, HCB, and mirex above human consumption tolerance limits set by the U.S. Food and Drug

Address correspondence to L.M. Schell, A&S 237, University at Albany, State University of New York, 1400 Washington Ave., Albany, NY 12222 USA. Telephone: (518) 442-5761. Fax: (518) 442-4563. E-mail: l.schell@albany.edu

We thank the Akwesasne Mohawk community for their cooperation and participation in this research.

This research was supported by grants from the National Institute of Environmental Health Sciences (ESO4913-10; ES10904-06) and the National Center on Minority Health and Health Disparities (5R01MD001120).

L.M.S. and A.P.D. have received financial compensation as experts in litigation involving human exposure to environmental contaminants at Akwesasne. D.O.C. has served as an expert witness in legal cases related to health effects of PCB exposure, with all reimbursement going to the Research Foundation of the University at Albany. The other authors declare they have no competing financial interests.

Received 22 May 2007; accepted 25 February 2008.

Administration (Forti et al. 1995; Sloan and Jock 1990). In 1986 and 1987, advisories against eating locally caught fish were issued. Local informants have attested to a reduction in consumption of locally caught fish, and studies of PCB levels in breast milk reported a decrease in levels subsequently that is consistent with adherence to the advisories (Fitzgerald et al. 2001). Thus, adolescents born after the fish advisories may have had less postnatal exposure than those born before the advisories, especially exposure to persistent PCBs.

Human health studies at Akwesasne, including an investigation of adolescent development, were prompted by the history of local environmental pollution from the neighboring industrial point sources. Long-standing reliance on locally caught fish and game (prime pathways of exposure) and concerns of community members about the health effects of environmental pollutants for themselves and future generations resulted in a community-based research collaboration.

Participants. The study began in 1995 and ended in 2000. Study methods, including recruitment and data collection protocols, laboratory analysis methods, and substitution methods for toxicant levels below the laboratory minimum detection limits, are briefly described here [for greater detail, see Schell et al. (2003)]. Akwesasne community members collected all data without prior knowledge of the exposure status of participants. The Institutional Review Board at the University at Albany, State University of New York, approved all study protocols. Informed consent, and assent from minors, was obtained from all participants.

The target population was defined as residents of Mohawk households located in the Akwesasne Mohawk Nation and in neighboring communities within 10 miles of Akwesasne. Mother–youth participant pairs were eligible for the study if they lived in the same household, and if the adolescent was between the ages of 10 and 16.99 years, not a twin, not diagnosed with a psychological or physical impairment, and not diagnosed with fetal alcohol syndrome or effects. Of 294 mother–youth pairs who met the study eligibility requirements and enrolled in the study, 271 continued participation and had blood available for analysis. Of the 271, 19 pairs had missing data for covariates included in the subsequent multiple regression analysis. Of the 252 remaining participants, 7 youth were missing either T_4 or FT_4 results. Of those remaining, 13 additional participants were excluded from this analysis because of a change in laboratory methods for the analysis of TSH and thyroid hormones, for a final sample size of 232.

Blood collection and laboratory analysis. Fasting blood specimens were collected at first rising by trained Mohawk staff and provided material for analysis of six toxicants (lead,

mercury, PCBs, p,p' -DDE, HCB, and mirex), cholesterol, triglycerides, TSH, and thyroid hormones. All samples were drawn between 1996 and 2000.

PCB and organochlorine pesticide analyses were conducted at the Exposure Assessment Laboratory of the University at Albany. High-resolution, ultratrace, congener-specific analysis was performed by parallel dual-column (splitless injection) gas chromatography with electron capture detection on an Agilent (Santa Clara, CA) 5890 instrument (DeCaprio et al. 2000). This method quantitates up to 83 individual PCB congeners and 18 PCB congeners as pairs or triplets, as well as p,p' -DDE, HCB, and mirex. Data were expressed on a whole-weight basis. Analyses of lead and mercury were conducted by Le Centre de Toxicologie due Quebec in Sainte-Foy, Quebec, Canada. Mercury analysis was based on cold-vapor atomic absorption spectrometry using a Pharmacia (Stockholm, Sweden) model 100 mercury monitor. Levels are reported as the sum of organic and inorganic mercury in micrograms per deciliter. Lead was analyzed by Zeeman-corrected graphite furnace atomic absorption spectrometry on a PerkinElmer (Waltham, MA) model 4100ZL instrument.

Assessment of cholesterol, triglycerides, TSH, T_3 , T_4 , and FT_4 was performed at the Clinical Chemistry and Hematology Laboratory, Wadsworth Center for Laboratories and Research, New York State Department of Health (Albany, NY). The facility is approved by the Clinical Laboratory Improvement Amendments and is a member of the Centers for Disease Control and Prevention (CDC) reference laboratory network for lipid measurements (Myers et al. 2000). Serum lipid concentrations were measured on a Hitachi 911 analyzer (Roche Diagnostics, Indianapolis, IN) using a cholesterol esterase and oxidase/peroxidase method for total cholesterol (Allain et al. 1974) and a glycerol kinase-based procedure that corrects for free glycerol in the specimen (Kohlmeier 1986) for triglycerides. Thyroid hormones were analyzed by ultrasensitive radioimmunoassay using standard methodologies (National Academy of Clinical Biology 2002). Sensitivity for TSH was 0.02 μ IU/mL, with a coefficient of variation of < 20%. Reference ranges for clinically normal values were 4.5–12.5 μ g/dL and 0.71–1.85 ng/dL for T_4 and FT_4 , respectively, 85–190 ng/dL for T_3 , and 0.3–5.0 for TSH.

Pollutants. Values below the method detection limits for lead, mercury, PCBs, and HCB were imputed using the U.S. Environmental Protection Agency recommended method (U.S. EPA 1998) for toxicants with rates of detection of \geq 50%. This method imputes a value for each datum below

the method detection limit based on the method detection limit value, the percentage of observations below the method detection limit, and the mean and variance of the detected observations. No imputation method was used for p,p' -DDE because all participants had detectable levels. To correct for skewness and normalize the distributions, lead, mercury, PCBs, p,p' -DDE, and HCB were natural log transformed. The 16 PCB congeners detected in > 50% of the sample are included in our analyses for hypothesis testing. PCB congeners were considered individually and in three groups: *a*) all 16 PCB congeners: PCB congeners detected in > 50% of the sample (PCB-50%): International Union of Pure and Applied Chemistry (IUPAC) congeners 52, 70, 74, 84, 87, 95, 99, 101[+ 90], 105, 110, 118, 149[+ 123], 138[+ 164 + 163], 153, 180, 187; *b*) eight persistent PCB congeners (PCB-PER8): IUPAC congeners 74, 99, 105, 118, 138[+ 164 + 163], 153, 180, 187; and *c*) eight nonpersistent congeners (PCB-NON8): IUPAC congeners 101[+ 90], 110, 95, 52, 149[+ 123], 84, 70, 87. Persistent PCBs are congeners with long (years) physiologic half-lives in humans (Brown 1994; Hansen 1998). IUPAC congeners 70 and 87 are classified as nonpersistent based on data presented by Brown (1994). Mirex levels were categorized into three groups because > 50% of the sample had levels below the method detection limit of 0.02 ppb: nondetects (< 0.02 ppb; 54.7%), low detects (0.02–0.03 ppb; 17.2% of the sample), and high detects (0.04–1.17 ppb; 28.0% of the sample).

Other variables. Additional information was obtained by interview with the mother of the youth, including sociodemographic variables and breast-feeding history, which was recorded as “any” or “none.” If a blood sample could not be analyzed at the laboratory, it was redrawn at a later point, occasionally creating a lag between collection of interview data and blood draw. The variable—time to blood analysis—describes this time difference. However, all biologic material analyzed to determine thyroid hormones and toxicant levels was from blood drawn at a single point in time.

Statistical analysis. We used multiple regression analysis to examine the effect of each toxicant on thyroid hormones when controlling for all other toxicants, as well as sex, age, triglycerides, cholesterol, breast-feeding, time of day when blood was collected, and duration of time between interview and blood draw. Covariates were chosen on the basis of bivariate associations ($p < 0.1$, t -tests, and correlations) with thyroid hormones and/or PCBs. A PCB-by-breast-feeding interaction was also included in multiple regression analyses. Breast-feeding can be a major route of postnatal PCB exposure, and levels of some PCB congeners typically differ by

breast-feeding status. Intercorrelations among toxicants were assessed because high intercorrelations can render the interpretation of multiple regression coefficients problematic.

Results

The mean (\pm SD) of adolescents' age was 13.3 ± 1.94 years. The mean triglyceride level was 86.9 mg/dL, and seven participants had triglyceride values over the laboratory reference range of 200 mg/dL. The mean cholesterol measure was 158.5 mg/dL, with 20 participants above the reference range of 200 mg/dL. The mean time of blood collection was 0831 ± 0101 hours, just 1 min earlier than the median time. The median time between interview and venipuncture was 1.1 days (-0.003 years), but some cases (27 of 232) had blood redrawn well after interview

to repeat the laboratory analysis. Thus, there is a difference between mean and median values for this variable. Nearly half (45.7%) of the youth were breast-fed as infants.

Mean levels of T_4 , FT_4 , T_3 , and TSH were within the laboratory reference range (Table 1). Eight participants had TSH levels above the laboratory reference range, and four had T_3 levels above the normal range. Few participants had T_3 , T_4 , or TSH levels below the laboratory reference range ($n = 3, 3$, and 4 , respectively). FT_4 and T_4 were significantly lower in youth who had been breast-fed, whereas TSH was significantly higher. T_3 did not differ.

Lead, mercury, HCB, p,p' -DDE, and PCB levels of Mohawk adolescents are described in Table 2. Toxicant levels in the study sample were consistent with a pattern of chronic exposure to multiple toxicants. The highest level of

lead in the sample was less than half of the CDC action level of 10 μ g/dL (CDC 1991). Mercury levels of Mohawk adolescents were at or below background levels of 0.1 – 0.8 μ g/dL that have been reported for the general population (Agency for Toxic Substances and Disease Registry 1999). Only one Mohawk adolescent had a mercury level ≥ 5.8 μ g/L, the blood mercury level equivalent to the U.S. EPA reference dose, compared with 5.66% of women of childbearing age in the general population (CDC 2004). PCB levels of adolescents and the proportion of persistent to nonpersistent congeners were consistent with both cumulative and recent exposure [58% of the most common congeners (PCB-50%) were persistent ones] (DeCaprio et al. 2005; Schell et al. 2003).

PCB-50%, PCB-PER8, and p,p' -DDE were significantly higher in the breast-fed participants (Table 2). When individual PCB congener levels were compared, all persistent congeners, except for PCB-105, had significantly higher values among adolescents who were breast-fed as infants. Levels of all non-persistent congeners, except for PCB-87, were not significantly different by breast-feeding status. No difference in lead, mercury, or HCB was observed by breast-feeding status. We also considered other factors that could differ between non-breast-fed and breast-fed youth (age, sex, two indices of socioeconomic status, triglycerides, cholesterol, and the youth's weight, height, and body mass index) and found no significant differences.

We used multivariate regression analyses to examine the effects of different measures of PCB exposure on thyroid hormone and TSH levels while controlling for other toxicants (HCB, p,p' -DDE, mirex, lead, mercury) and additional covariates [age, sex, cholesterol, triglycerides, time of collection (time of day), time to blood analysis (years), breast-feeding, and breast-feeding by PCB interaction]. Results are presented for two regression analyses using PCB-PER8 as the PCB measure to predict log-transformed TSH (Table 3) and FT_4 (Table 4). PCB-PER8 was positively associated with TSH. A breast-feeding-by-PCB-PER8 interaction was statistically significant. From this regression analysis, we calculated the effect of PCB-PER8 on TSH and FT_4 in both the breast-fed and non-breast-fed adolescents while holding all other model variables constant at their respective means. As PCB-PER8 levels rise from 0.204 ppb at the 5th percentile to 0.871 ppb at the 95th percentile, estimated TSH levels increase by 1.51 μ U/mL among adolescents who were not breast-fed as infants. Among adolescents who were breast-fed as infants, TSH levels are not significantly related to PCB-PER8; at the 95th percentile of PCB-PER8, the estimated TSH level is 2.8 μ U/mL compared with 3 μ U/mL at the 50th percentile level. Repeating the analyses without the individuals

Table 1. Thyroid hormone levels among Akwesasne Mohawk youth ($n = 232$).

Hormones	Reference range ^a	Mean \pm SD	Percent outside range	
			Below	Above
T_4	4.5–12.5 μ g/dL	7.1 \pm 1.34	1.3	0
FT_4	0.71–1.85 ng/dL	1.1 \pm 0.16	0	0
T_3	85–190 ng/dL	138.1 \pm 25.93	1.3	1.7
TSH	0.3–5.0 μ U/mL	2.5 \pm 1.46	1.7	3.5

^aReference range provided by the Clinical Chemistry and Hematology Laboratory, Wadsworth Center for Laboratories and Research, New York State Department of Health (unpublished data).

Table 2. Toxicant levels of Akwesasne Mohawk adolescents: breast-fed and non-breast-fed ($n = 232$).

Toxicant	Non-breast-fed ($n = 126$)				Breast-fed ($n = 106$)				<i>t</i>
	GM	Median	SD	Max	GM	Median	SD	Max	
Total PCBs (ppb)									
Zero substitution ^a	0.71	0.68	0.668	3.48	0.95	0.95	0.806	4.52	–2.7**
MDL substitution ^b	1.53	1.41	0.557	3.79	1.75	1.67	0.695	4.74	–2.9**
MDL substitution ^c	2.27	2.14	0.452	4.16	2.47	2.39	0.589	4.96	–3.1**
PCB-50% (ppb) ^{d,e}	0.59	0.55	0.272	1.66	0.77	0.77	0.405	2.36	–4.7**
PCB-PER8 (ppb) ^{d,f}	0.34	0.32	0.168	1.44	0.47	0.46	0.259	1.38	–6.0**
PCB-NON8 (ppb) ^{d,g}	0.25	0.23	0.148	0.94	0.28	0.25	0.203	1.33	–1.7
Persistent PCB congeners (ppb) ^d									
74	0.02	0.02	0.010	0.09	0.03	0.03	0.029	0.22	–5.9**
99	0.04	0.04	0.021	0.13	0.05	0.06	0.034	0.21	–4.3**
105	0.02	0.02	0.011	0.06	0.02	0.02	0.019	0.13	–1.6
118	0.06	0.06	0.029	0.16	0.07	0.07	0.050	0.28	–3.2**
138[+ 163 = 164]	0.06	0.07	0.041	0.34	0.08	0.08	0.056	0.28	–3.5**
153	0.07	0.07	0.047	0.39	0.11	0.11	0.107	0.98	–5.7**
180	0.03	0.03	0.038	0.27	0.05	0.05	0.054	0.35	–4.9**
187	0.02	0.01	0.013	0.13	0.02	0.02	0.014	0.08	–4.3**
Nonpersistent PCB congeners (ppb) ^d									
52	0.03	0.02	0.027	0.16	0.03	0.02	0.029	0.14	–0.7
70	0.02	0.02	0.015	0.11	0.02	0.02	0.019	0.08	–1.5
84	0.02	0.02	0.007	0.05	0.02	0.02	0.010	0.08	–1.3
87	0.03	0.04	0.021	0.13	0.04	0.04	0.027	0.16	–2.3*
95	0.02	0.02	0.014	0.08	0.03	0.02	0.023	0.15	–1.5
101[+ 90]	0.05	0.04	0.034	0.20	0.05	0.05	0.048	0.30	–1.6
110	0.05	0.05	0.033	0.16	0.06	0.06	0.049	0.34	–1.6
149[+ 123]	0.02	0.02	0.016	0.08	0.02	0.02	0.021	0.12	–0.7
p,p' -DDE (ppb)	0.31	0.31	0.163	1.23	0.45	0.42	0.416	2.93	–5.9**
HCB (ppb) ^d	0.03	0.03	0.022	0.19	0.04	0.04	0.017	0.13	–1.6
Blood lead (μ g/dL) ^d	0.80	1.30	0.965	4.80	0.76	1.45	0.907	3.50	0.3
Mercury (μ g/dL) ^d	0.09	0.10	0.093	0.58	0.09	0.08	0.096	0.52	0.6

Abbreviations: GM, geometric mean; Max, maximum; MDL, method detection limit.

^aValues below the MDL were substituted with zero. ^bValues below the MDL were substituted with one half the MDL.

^cValues below the MDL were substituted with the MDL. ^dValues below the MDL were substituted with values calculated following the U.S. EPA recommended method. ^eCongeners with $\geq 50\%$ detection rate. IUPAC congeners: 52, 70, 74, 84, 87, 95, 99, 101[+90], 105, 110, 118, 149[+ 123], 138[+ 164 + 163], 153, 180, 187. ^fSum of eight persistent PCB congeners with $\geq 50\%$ detection rate. IUPAC congeners: 74, 99, 105, 118, 138[+ 164 + 163], 153, 180, 187. ^gSum of eight nonpersistent PCB congeners with $\geq 50\%$ detection rate. IUPAC congeners: 52, 70, 84, 87, 95, 101[+ 90], 110, 149[+ 123]. * $p < 0.05$. ** $p < 0.01$.

with TSH, T_4 , T_3 levels outside the laboratory range produced results, including those for the interaction effect, that were unchanged for effects of PCB-PER8, and only slightly different for effects of PCB-50% and lead. The beta coefficient estimating the relationship between PCB50% and T_4 increased trivially from -0.06 to -0.065, with a change in p -value from 0.057 to 0.033. The effect of lead in the model was attenuated, and p -values increased to just above 0.05.

Table 4 shows a negative effect of PCB-PER8 on FT_4 . The PCB-PER8 by breast-feeding interaction term was positive but not statistically significant. Applying this model to the non-breast-fed group, predicted FT_4 levels decrease from 3.1 ng/dL at the 5th percentile of PCB-PER8 to 2.9 ng/dL at the 50th percentile and to 2.7 ng/dL at the 95th percentile. Among breast-fed adolescents, FT_4 was essentially unchanged with increasing PCB-PER8.

PCB-NON8 was not significantly related to TSH but was significantly and negatively related to FT_4 . Because congener groups were related to either TSH, FT_4 , or both, the constituent congeners were tested individually (Table 5). Of the persistent congeners, 118, 138[+ 163 + 164], and 153 were positively associated with TSH levels, and only one nonpersistent congener (110) showed a relationship with TSH. Two persistent and six nonpersistent PCB congeners were negatively associated with FT_4 , as were all three summary PCB measures.

HCB was negatively associated with T_4 , and lead was positively associated with T_3 (results not shown in table). Results for HCB and lead were consistent regardless of which measure of PCB burden was included in the multivariate model.

On examination of the interrelationships among toxicants, p,p' -DDE was found to be positively correlated with PCB-50%, PCB-PER8, and HCB, with coefficients of 0.46, 0.60, and 0.39, respectively. Collinearity among toxicants was not considered an analytical complication, because the largest coefficient was well below 0.80. To ensure that collinearity was not influential, the regression analysis was performed with models trimmed of covariates that were not significant ($p < 0.1$) in any single model (time of blood collection, p,p' -DDE, mirex nondetects vs. detects). Trimmed models were little changed (< 20% change in beta coefficients, and the same PCB and thyroid effects were significant at $p < 0.05$; results not shown) when compared with full models that included time of blood collection, p,p' -DDE, and mirex nondetects versus detects.

Discussion

The most striking observation in this study is the demonstration of a significant positive relationship between serum PCB levels and

TSH in adolescents who were not breast-fed, and the lack of such a relationship in adolescents who were breast-fed—despite the higher PCB levels found in breast-fed adolescents. There was also a negative but less significant relationship between serum PCB levels and FT_4 in non-breast-fed, but not in breast-fed, adolescents.

Breast-feeding has been shown to be the largest postnatal source of PCB burden in chronically exposed populations (Chao et al. 2004; Lackman et al. 2004; Patandin et al. 1999). One would therefore expect PCB effects to be greater in those youth with higher PCB levels, but this is not the case here. The non-breast-fed youth displayed evidence of stronger associations between PCB levels and TSH and FT_4 than the breast-fed group with higher PCB levels. It is possible that breast-feeding has a beneficial effect, reducing the impact of prenatal PCB exposure on measures of thyroid function. Studies have documented the benefits of breast-feeding on development, mortality, and morbidity (Anderson et al.

1999; Howie et al. 1990; Kramer et al. 2001; Singhal et al. 2001; WHO Collaborative Study Team on the Role of Breastfeeding on the Prevention of Infant Mortality 2000; Wilson et al. 1998). However, the suggestion that breast-feeding is protective of PCB-induced alterations of thyroid function is not concordant with the higher levels of PCBs in breast-fed adolescents and the expectation of a dose-response relationship.

A more likely explanation for our observations is that prenatal exposure to PCBs alters thyroid function to a greater degree than early postnatal exposure or exposure during childhood and adolescence. The notion of a critical window is consistent with findings on organochlorine exposure and brain development. In newborn mice where brain development is in a critical period, exposure to persistent organochlorines, such as PCBs, has been shown to cause severe, sometimes irreversible, brain disruption (Eriksson 1997; Eriksson et al. 2002). If the effect of PCBs on thyroid function is sensitive to the timing of

Table 3. Multivariate regression analysis predicting TSH levels (μ U/mL)^a of Akwesasne Mohawk adolescents ($n = 232$).

Characteristic	β	SE	Std β	t	p -Value
Constant	1.062	0.674		1.576	0.117
Age (years)	0.026	0.021	0.082	1.250	0.213
Sex (male/female)	-0.187	0.080	-0.150	-2.326	0.021
Breast-feeding (any/none)	-0.202	0.203	-0.162	-0.993	0.322
Time of blood collection	-0.000	0.000	-0.015	-0.230	0.818
Time to blood analysis (years)	-0.065	0.040	-0.103	-1.639	0.103
Cholesterol (mg/dL)	-0.001	0.002	-0.055	-0.778	0.438
Triglycerides (mg/dL)	0.003	0.001	0.234	3.323	0.001
HCB (ppb) ^a	0.084	0.106	0.055	0.798	0.426
p,p' -DDE (ppb) ^a	-0.076	0.109	-0.060	-0.695	0.488
Blood lead (μ g/dL) ^a	-0.017	0.029	-0.036	-0.562	0.575
Mercury (μ g/dL) ^a	-0.026	0.053	-0.031	-0.485	0.628
Mirex (nondetects vs. detects)	-0.104	0.108	-0.083	-0.960	0.338
Mirex (high detects vs. low and nondetects)	-0.219	0.119	-0.158	-1.840	0.067
PCB-PER8 (ppb) ^a	0.431	0.149	0.309	2.899	0.004
PCB-PER8 \times breast-feeding	-0.505	0.192	-0.396	-2.626	0.009

Std, standardized.

^aNatural log transformed.

Table 4. Multivariate regression analysis predicting FT_4 levels (ng/dL) of Akwesasne Mohawk adolescents ($n = 232$).

Characteristic	β	SE	Std β	t	p -Value
Constant	1.009	0.183		5.501	0.000
Age (years)	-0.007	0.006	-0.085	-1.237	0.217
Sex (male/female)	0.009	0.022	0.028	0.414	0.680
Breast-feeding (any/none)	0.025	0.055	0.077	0.451	0.652
Time of blood collection	-0.000	0.000	-0.052	-0.786	0.433
Time to blood analysis (years)	0.019	0.011	0.118	1.793	0.074
Cholesterol (mg/dL)	0.000	0.000	0.085	1.156	0.249
Triglycerides (mg/dL)	0.000	0.000	-0.068	-0.927	0.355
HCB (ppb) ^a	-0.027	0.029	-0.067	-0.935	0.351
p,p' -DDE (ppb) ^a	-0.003	0.030	-0.008	-0.092	0.926
Blood lead (μ g/dL) ^a	0.001	0.008	0.010	0.145	0.885
Mercury (μ g/dL) ^a	0.007	0.014	0.033	0.501	0.617
Mirex (nondetects vs. detects)	-0.010	0.029	-0.030	-0.330	0.741
Mirex (high detects vs. low and nondetects)	-0.016	0.032	-0.046	-0.508	0.612
PCB-PER8 (ppb) ^a	-0.099	0.040	-0.272	-2.443	0.015
PCB-PER8 \times breast-feeding	0.066	0.052	0.198	1.262	0.208

Std, standardized.

^aNatural log transformed.

the exposure also, an effect of PCBs that occurs only with prenatal exposure might be observable only among adolescents who experienced little postnatal exposure to PCBs. For non-breast-fed adolescents, PCB levels would be expected to more closely reflect prenatal exposure. Breast-feeding exposes children to a comparatively large dose of PCBs and is a major source of overall PCB body burden (Ayotte et al. 2003; Lackman et al. 2004; Lanting et al. 1998). Thus, breast-feeding may add PCBs that obscure the relationship between prenatal PCB exposure and thyroid function by adding random variation in PCB levels, rather than protecting the thyroid from the disruptive influence of PCBs. Breast-feeding is not associated with additional risk to thyroid disruption, despite the higher levels of PCBs resulting from breast-feeding.

There are a number of reasons to suspect that prenatal PCB exposure might alter thyroid function later in life. PCBs disturb differentiation of normal human neural progenitor cells, a thyroid hormone-dependent process (Fritsche et al. 2005); inhibit thyroid hormone-dependent extension of dendrites (Kimura-Kuroda et al. 2005); and cause a decrease in pituitary and thyroid responses to thyrotropin-releasing hormone stimulation (Khan and Hansen 2003). Gauger et al. (2004) have reported that PCBs have direct actions on several thyroid hormone-responsive genes in the fetal rat brain and increase the expression of neuroendocrine-specific protein A, RC3/neurogranin, and Oct-1. These actions occur independently of the

reduced circulating levels of T_3 and T_4 in the dam. Miyazaki et al. (2004) found that PCBs suppress thyroid receptor-mediated transcription and suggested that this is particularly the case in the developing nervous system. Thus, there are several possible mechanisms whereby PCB exposure during development might alter thyroid function permanently.

Prenatal exposure to PCBs also causes a greater and more persistent alteration of other organ systems than does postnatal exposure. In children, decrements in neurobehavioral function resulting from PCB and dioxin exposure are primarily a result of prenatal exposure (Lai et al. 2001; Vreugdenhil et al. 2002). Prenatal exposure to PCBs and dibenzofurans has been found to alter semen quality (Guo et al. 2000) and sperm function (Hsu et al. 2003) in adult humans and alter fertility in adult rats (Kuriyama and Chahoud 2004).

Our results show that not all PCB congeners have similar effects. We observed statistically significant, positive associations between TSH levels and PCB congeners 110, 118, 138[+ 163 + 164], 153, as well as with two PCB groupings (PCB-50%, PCB-PER8). Of the four individual congeners/triplets, three are persistent, three are di-*ortho* congeners, and all four are highly chlorinated. Negative associations were observed between FT_4 levels and three congener groupings (PCB-50%, PCB-PER8, PCB-NON8), as well as congeners 52, 70, 84, 87, 101[+ 90], 138[+ 163 + 164], 149[+ 123], and 153. Six of these eight congeners are highly chlorinated, and five are

di-*ortho* congeners. Only congener 87 was associated with T_4 levels, and none of the tested PCB congeners or groupings was associated with T_3 levels. The results presented here on the relationships of PCBs to TSH, FT_4 , and T_3 are consistent with a preliminary report on a sample of 113 Mohawk adolescents who participated in this study (Schell et al. 2002), and with a recent report showing an increasing incidence rate of hypothyroidism among patients using the St. Regis Mohawk Health Service between 1992 and 1995 (Negoiita et al. 2001).

The positive association observed here between PCBs and TSH, and the negative association between PCBs and thyroid hormones is supported by the literature. PCBs (as individual congeners or mixtures) have repeatedly been shown to alter thyroid function and hormone levels in experimental animals, including thyroid-hormone suppression and cell-mediated immunomodulation (Brouwer et al. 1998; Hallgren et al. 2001; Kato et al. 1998; Kuriyama et al. 2003; Li and Hansen 1996a, 1996b; McNabb and Fox 2003; Morse et al. 1993; Ness et al. 1993; Seo et al. 1995; Smits et al. 2002). *In utero* exposure of rats produces depressed plasma T_4 levels in late gestation and in newborns (Morse et al. 1993; Ness et al. 1993; Seo et al. 1995), and prepubertal exposure can depress serum T_4 levels in rats as well (Li and Hansen 1996a, 1996b). Studies of PCB-thyroid effects in humans have been less consistent. In men and women who consumed fish from the Great Lakes, serum PCBs were associated with lower T_4 levels. Inconsistent associations were found with TSH and PCBs (Persky et al. 2001). Using data from the National Health and Nutrition Examination Survey (NHANES) (1999–2002), Turyk et al. (2007) found PCBs to be positively related to TSH among older women, yet inversely associated among older men. Additionally, a negative relationship of TEQ to T_4 was seen in both men and women, with a stronger correlation among women. No associations were reported for younger NHANES participants (Turyk et al. 2007). A study of men between 20 and 64 years of age found inverse relationships between T_3 and two persistent PCB congeners (138, 153), the sum of PCBs, and HCB (Meeker et al. 2007). In the Dutch multicenter study of newborns, higher PCB and dioxin levels in breast milk were positively associated with TSH levels and negatively associated with FT_4 levels (Koopman-Esseboom et al. 1994; Sauer et al. 1994). One-year-old Japanese infants who were breast-fed had lower T_3 and T_4 levels at higher levels of polychlorinated dibenzo-*p*-dioxins and dibenzofurans and coplanar PCBs in mothers' milk, although TSH levels were unrelated (Nagayama et al. 1998). Positive associations between congener 118 and TSH levels have been reported in 320 German schoolchildren

Table 5. Association between different measures of PCBs, each entered individually in multiple regression analyses,^a and thyroid hormones and TSH ($n = 232$).

PCBs (ppb) ^{b,c}	Standardized β			
	TSH ^b	FT_4	T_4	T_3
PCB-50% ^d	0.238*	-0.269**	-0.134	-0.053
PCB-PER8 ^e	0.309**	-0.272*	-0.121	-0.020
PCB-NON8 ^f	0.131	-0.242*	-0.133	-0.061
Persistent PCB IUPAC congeners				
74	0.145	-0.229	-0.111	0.005
99	0.117	-0.173	-0.007	-0.011
105	0.014	-0.022	-0.040	-0.047
118	0.267**	-0.191	-0.049	-0.026
138[+ 163 + 164]	0.349**	-0.234*	-0.116	-0.119
153	0.285**	-0.324**	-0.158	-0.037
180	0.193	-0.125	-0.067	0.037
187	0.164	-0.124	-0.078	-0.033
Nonpersistent PCB IUPAC congeners				
52	0.056	-0.191*	-0.083	-0.100
70	0.121	-0.236*	-0.119	-0.091
84	0.062	-0.226*	-0.120	-0.011
87	0.150	-0.267**	-0.189*	-0.028
95	0.014	-0.196	-0.141	-0.055
101[+ 90]	0.103	-0.214*	-0.095	-0.014
110	0.186*	-0.167	-0.111	-0.070
149[+ 123]	0.176	-0.248*	-0.073	-0.018

^aControlling for age, sex, time of day of collection, time to blood analysis, breast-feeding (any/none), cholesterol, triglycerides, breast-feeding \times PCBs, lead, mercury, HCB, DDE, and mirex. ^bLog transformed. ^cValues below the method detection limit were substituted with values calculated following the U.S. EPA recommended method. ^dCongeners with $\geq 50\%$ detection rate. IUPAC congeners: 52, 70, 74, 84, 87, 95, 99, 101[+ 90], 105, 110, 118, 149[+ 123], 138[+164+163], 153, 180, 187. ^eSum of eight persistent PCB congeners with $\geq 50\%$ detection rate. IUPAC congeners: 74, 99, 105, 118, 138[+164+163], 153, 180, 187. ^fSum of eight nonpersistent PCB congeners with $\geq 50\%$ detection rate. IUPAC congeners: 101[+ 90], 110, 95, 52, 149[+123], 84, 70, 87. * $p < 0.05$. ** $p < 0.01$.

between 7 and 10 years of age (Osius et al. 1999), and 98 newborns in Spain (Ribas-Fito et al. 2003), as well as in the present study. A longitudinal study of 38 breast-fed infants in the Netherlands reported a positive association between dioxin concentration in breast milk and TSH levels at 11 weeks (Pluim et al. 1992, 1993). However, Pluim and coworkers (1992, 1993) also observed higher T_4 levels at birth, 1 week, and 11 weeks, contrary to results reported here as well as by others described above. Negative associations between serum PCB levels and FT_4 have been reported in neonates from coastal communities in Quebec (Sandau et al. 2002) and adults from a Spanish village (Sala et al. 2001), whereas in an adult population from Quebec there was a negative relationship between levels of three PCB congeners (138, 153, and 180) and T_3 but not with TSH or FT_4 (Takser et al. 2005).

PCBs have long been suspected to affect thyroid hormone signaling because of structural similarities between PCBs and thyroid hormones (Porterfield 1994). Evidence suggests that two plausible mechanisms may be responsible for the PCB-induced reduction in thyroid hormones, particularly T_4 : disruption of thyroid hormone transport (Cheek et al. 1999; Darnerud et al. 1996) and induction of hepatic metabolism (Morse et al. 1993; Zhou et al. 2001). Results presented here are consistent with either mechanism of action. In the former mechanism, thyroid hormones are bound and transported primarily by two plasma proteins in humans, transthyretin (TTR) and thyroid-binding globulin (TBG) (Larsen et al. 2003). TTR is more likely to be involved as the transport protein in humans, because hydroxylated PCBs have greater affinities for TTR than does T_4 (Cheek et al. 1999; Lans et al. 1993, 1994; Rickenbacher et al. 1986). In contrast, few hydroxylated or unmetabolized PCBs bind TBG (Lans et al. 1993, 1994). In one study, after PCB exposure, the degree of thyroid hormone reduction in rats coincided with binding of T_4 to the plasma thyroid hormone transporter TTR, suggesting that PCB-induced T_4 reductions were attributed primarily to disturbed transport (Hallgren and Darnerud 2002). The second mechanism that has been investigated is induction of hepatic metabolism. Several studies have suggested that PCB-induced T_4 reductions result from increased metabolism of the hepatic microsomal enzyme uridine diphosphoglucuronosyl transferase (UDPGT). This induction is primarily the result of aryl hydrocarbon receptor activation by dioxin-like PCB congeners (Barter and Klaassen 1992, 1994; Beetstra et al. 1991; Morse et al. 1993; Van Birgelen et al. 1995; Zhou et al. 2001). UDPGT catalyzes glucuronidation of T_4 and consequently increases biliary excretion of T_4 (Bastomsky 1974). However, others suggest a

moderate or nonsignificant effect of UDPGT induction on PCB-mediated decreases in T_4 levels (Hallgren and Darnerud 2002; Hallgren et al. 2001). Because of insufficient specimen availability, we could not obtain data for the most potent dioxin-like congeners (CBs 126 and 169), which are typically present in human serum at levels two or three orders of magnitude lower than the most prevalent congeners. Therefore, we could not employ a dioxin-like TEQ approach (Van den et al. 2006) that may have provided additional support for the latter mechanism.

We also observed a negative association of HCB with T_4 and a positive association between lead and T_3 . The negative association of HCB with T_4 levels among Akwesasne adolescents is consistent with the literature. Numerous animal studies have demonstrated HCB-induced hypothyroidism, with T_4 levels being particularly sensitive in rats (Alvarez et al. 2005; Foster et al. 1993; Kleiman de Pisarev et al. 1990, 1995; Rozman et al. 1986; van Raaij et al. 1993). Lower T_4 levels associated with HCB exposure may be attributed to peripheral disposition of T_4 (Kleiman de Pisarev et al. 1989) or increased hepatic T_4 metabolism (Kleiman de Pisarev et al. 1990). It has also been suggested that HCB affects the thyroid in rats via its metabolites, particularly the main metabolite pentachlorophenol, an effective competitor for T_4 binding sites (van Raaij et al. 1991a, 1991b). In Turkey, 37% of patients with HCB-induced porphyria also developed enlarged thyroid glands (Gocmen et al. 1986). In a highly exposed, rural population residing in Catalonia, Spain, HCB levels were associated with lower T_4 levels (Sala et al. 2001). However, no relationship was observed between HCB and TSH levels in 98 newborns born in a highly HCB-polluted area (Ribas-Fito et al. 2003).

Although animal studies have shown lead to inhibit thyroid function and reduce circulating levels of T_3 and T_4 (Sandstead 1967; Shrivastava et al. 1987; Singh and Dhawan 1999), studies of thyroid functioning in humans have yielded mixed results. Some report no relationship (Erfurth et al. 2001; Gennart et al. 1992; Schumacher et al. 1998), whereas others report depressed thyroid hormone levels and/or function (Liang et al. 2003; Robins et al. 1983; Sandstead et al. 1969). It is also possible that lead may have different effects on thyroid hormones at different levels of exposure. In a study among occupationally exposed men, lead was positively associated with T_3 , T_4 , FT_4 , and TSH among workers with blood lead levels from 8 to 50 $\mu\text{g}/\text{dL}$ and negatively associated with T_3 and T_4 when blood lead levels exceeded 50 $\mu\text{g}/\text{dL}$ (Lopez et al. 2000). Directional differences in reported effects may stem from different levels and durations of exposure, and

may explain the positive association with T_3 observed here among Akwesasne adolescents.

Toxicant levels in Akwesasne youth (Schell et al. 2003) are somewhat lower than reported levels in older children and adolescents from several other studies (Karmaus et al. 2001; Mazhitova et al. 1998; Nawrot et al. 2002; Osius et al. 1999; Staessen et al. 2001). These populations with higher toxicant levels typically involve acute or known, well-defined sources of exposure. This suggests that toxicant body burdens reported here and, consequently, associations observed with thyroid hormones, may be present in other populations without large or acute exposures. In addition, it is unlikely that iodine insufficiency is responsible for these results. Sodium intake estimated by a semiquantitative food frequency questionnaire (Block et al. 1990, 1992; National Cancer Institute 1999) was more than sufficient to provide the recommended daily intake of iodine, assuming conservatively that only one third of the consumed sodium was iodized. An alternative explanation for thyroid effects in this sample is exposure to fluoride, a thyrotoxicant (National Research Council 2006). In 1980, cattle on Cornwall Island within the reservation and immediately downwind from an aluminum plant exhibited fluorosis despite a fluoride level in forage well below the tolerance level set by the National Academy of Sciences (Crissman et al. 1980). Human populations were exposed to fluoride through locally grown fruits and vegetables rather than from fish, the main route of PCB exposure. There was little evidence of effects on thyroid function: Men and women in the high fluoride group did not differ from those in a low exposure group in terms of T_4 ; high fluoride women had higher TSH, whereas the men had a significantly lower TSH level (Selikoff et al. 1983) than those in the low fluoride group.

In summary, our results demonstrate a reduction in thyroid function in adolescents in relation to their serum levels of PCBs, but demonstrate that this relationship is much stronger in adolescents who were not breast-fed, even though breast-fed adolescents have higher serum PCB levels. These observations are consistent with the hypothesis that prenatal exposure to PCBs alters thyroid function in a long-lasting manner.

REFERENCES

- Agency for Toxic Substances and Disease Registry. 1999. Toxicological Profile for Mercury. Atlanta, GA:U.S. Department of Health and Human Services, Public Health Service.
- Allain CC, Poon LS, Chan CS, Richmond W, Fu PC. 1974. Enzymatic determination of total serum cholesterol. *Clin Chem* 20:470–475.
- Alvarez L, Hernandez S, Martinez-de-Mena R, Kolliker-Frers R, Obregon MJ, Kleiman de Pisarev DL. 2005. The role of type I and type II 5' deiodinases on hexachlorobenzene-induced alteration of the hormonal thyroid status. *Toxicology* 207:349–362.

- Anderson JW, Johnstone BM, Remley DT. 1999. Breast-feeding and cognitive development: a meta-analysis. *Am J Clin Nutr* 70:525–535.
- Ayotte P, Muckle G, Jacobson JL, Jacobson SW, Dewailly É. 2003. Assessment of pre- and postnatal exposure to polychlorinated biphenyls: lessons from the Inuit Cohort Study. *Environ Health Perspect* 111:1253–1258.
- Barter RA, Klaassen CD. 1992. UDP-glucuronosyltransferase inducers reduce thyroid hormone levels in rats by an extrathyroidal mechanism. *Toxicol Appl Pharmacol* 113:36–42.
- Barter RA, Klaassen CD. 1994. Reduction of thyroid hormone levels and alteration of thyroid function by four representative UDP-glucuronosyltransferase inducers in rats. *Toxicol Appl Pharmacol* 128:9–17.
- Bastomsky CH. 1974. Effects of a polychlorinated biphenyl mixture (Aroclor 1254) and DDT on biliary thyroxine excretion in rats. *Endocrinology* 95:1150–1155.
- Beetstra JB, van Engelen JG, Karels P, van der Hoek HJ, de Jong M, Docter R, et al. 1991. Thyroxine and 3,3',5-triiodothyronine are glucuronidated in rat liver by different uridine diphosphate-glucuronyltransferases. *Endocrinology* 128:741–746.
- Block G, Hartman AM, Naughton D. 1990. A reduced dietary questionnaire: development and validation. *Epidemiology* 1:58–64.
- Block G, Thompson FE, Hartman AM, Larkin FA, Guire KE. 1992. Comparison of two dietary questionnaires validated against multiple dietary records collected during a 1-year period. *J Am Diet Assoc* 92:686–693.
- Brouwer A, Morse DC, Lans MC, Schuur AG, Murk AJ, Klasson-Wehler E, et al. 1998. Interactions of persistent environmental organohalogenes with the thyroid hormone system: mechanisms and possible consequences for animal and human health. *Toxicol Ind Health* 14:59–84.
- Brown JF Jr. 1994. Determination of PCB metabolic, excretion, and accumulation rates for use as indicators of biological response and relative risk. *Environ Sci Technol* 28:2295–2305.
- Brucker-Davis F. 1998. Effects of environmental synthetic chemicals on thyroid function. *Toxicol* 8:827–856.
- Carpenter DO. 2006. Polychlorinated biphenyls (PCBs): routes of exposure and effects on human health. *Rev Environ Health* 21:1–23.
- CDC. 1991. Preventing Lead Poisoning in Young Children. Atlanta, GA:Centers for Disease Control and Prevention.
- CDC (Centers for Disease Control and Prevention). 2004. Blood mercury levels in young children and childbearing-aged women—United States, 1999–2002. *MMWR* 53:1018–1020.
- Chao HR, Wang SL, Lee CC, Yu HY, Lu YK, Papke O. 2004. Level of polychlorinated dibenzo-*p*-dioxins, dibenzofurans and biphenyls (PCDD/Fs, PCBs) in human milk and the input to infant body burden. *Food Chem Toxicol* 42:1299–1308.
- Cheek AO, Kow K, Chen J, McLachlan JA. 1999. Potential mechanisms of thyroid disruption in humans: interaction of organochlorine compounds with thyroid receptor, transthyretin, and thyroid-binding globulin. *Environ Health Perspect* 107:273–278.
- Crissman JW, Maylin GA, Krook L. 1980. New York State and U.S. federal fluoride pollution standards do not protect cattle health. *Cornell Vet* 70:183–192.
- Darnerud PO, Morse D, Klasson-Wehler E, Brouwer A. 1996. Binding of a 3,3', 4,4'-tetrachlorobiphenyl (CB-77) metabolite to fetal transthyretin and effects on fetal thyroid hormone levels in mice. *Toxicology* 106:105–114.
- Daston GP, Cook JC, Kavlock RJ. 2003. Uncertainties for endocrine disruptors: our view on progress. *Toxicol Sci* 74:245–252.
- DeCaprio AP, Johnson GW, Tarbell AM, Carpenter DO, Chiarenzelli JR, Morse GS, et al. 2005. Polychlorinated biphenyl (PCB) exposure assessment by multivariate statistical analysis of serum congener profiles in an adult Native American population. *Environ Res* 98:284–302.
- DeCaprio AP, Tarbell AM, Bott A, Wagemaker DL, Williams RL, O'Hehir CM. 2000. Routine analysis of 101 polychlorinated biphenyl congeners in human serum by parallel dual-column gas chromatography with electron capture detection. *J Anal Toxicol* 24:403–420.
- Erfurth EM, Gerhardsson L, Nilsson A, Rylander L, Schutz A, Skerfving S, et al. 2001. Effects of lead on the endocrine system in lead smelter workers. *Arch Environ Health* 56:449–455.
- Eriksson P. 1997. Developmental neurotoxicity of environmental agents in the neonate. *Neurotoxicology* 18:719–726.
- Eriksson P, Viberg H, Jakobsson E, Örn U, Fredriksson A. 2002. A brominated flame retardant, 2,2',4,4',5-pentabromodiphenyl ether: uptake, retention, and induction of neurobehavioral alterations in mice during a critical phase of neonatal brain development. *Toxicol Sci* 67:98–103.
- Fitzgerald EF, Hwang S-A, Deres DA, Bush B, Cook K, Worswick P. 2001. The association between local fish consumption and DDE, mirex, and HCB concentrations in the breast milk of Mohawk women at Akwesasne. *J Expo Anal Environ Epidemiol* 11:381–388.
- Forti A, Bogdan KG, Horn E. 1995. Health Risk Assessment for the Akwesasne Mohawk Population from Exposure to Chemical Contaminants in Fish and Wildlife. Albany, NY:New York State Department of Health.
- Foster WG, Pentick JA, McMahon A, Lecavalier PR. 1993. Body distribution and endocrine toxicity of hexachlorobenzene (HCB) in the female rat. *J Appl Toxicol* 13:79–83.
- Fritsche E, Cline JE, Nguyen NH, Scanlan TS, Abel J. 2005. Polychlorinated biphenyls disturb differentiation of normal human neural progenitor cells: clue for involvement of thyroid hormone receptors. *Environ Health Perspect* 113:871–876.
- Gauger KJ, Kato Y, Haraguchi K, Lehmler HJ, Robertson LW, Bansal R, et al. 2004. Polychlorinated biphenyls (PCBs) exert thyroid hormone-like effects in the fetal rat brain but do not bind to thyroid hormone receptors. *Environ Health Perspect* 112:516–523.
- Gennart J-P, Bernard A, Lauwerys R. 1992. Assessment of thyroid, testes, kidney and autonomic nervous system function in lead-exposed workers. *Int Arch Occup Environ Health* 64:49–57.
- Gocmen A, Peters HA, Cripps DJ, Morris CR, Dogramaci I. 1986. Porphyria turcica: hexachlorobenzene-induced porphyria. *IARC Sci Publ* 77:567–573.
- Guo YL, Ping-Chi H, Chao-Chin H, Lambert GH. 2000. Semen quality after prenatal exposure to polychlorinated biphenyls and dibenzofurans. *Lancet* 356:1240–1241.
- Hagmar L. 2003. Polychlorinated biphenyls and thyroid status in humans: a review. *Thyroid* 13:1021–1028.
- Hallgren S, Darnerud PO. 2002. Polybrominated diphenyl ethers (PBDEs), polychlorinated biphenyls (PCBs) and chlorinated paraffins (CPs) in rats—testing interactions and mechanisms for thyroid hormone effects. *Toxicology* 177:227–243.
- Hallgren S, Sinjari T, Hakansson H, Darnerud PO. 2001. Effects of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) on thyroid hormone and vitamin A levels in rats and mice. *Arch Toxicol* 75:200–208.
- Hansen LG. 1998. Stepping backward to improve assessment of PCB congener toxicities. *Environ Health Perspect* 106:171–189.
- Howie PW, Forsyth JS, Ogston SA, Clark A, Florey CD. 1990. Protective effect of breast feeding against infection. *Br Med J* 300:11–16.
- Hsu PC, Huang W, Yao WJ, Wu MH, Guo YL, Lambert GH. 2003. Sperm changes in men exposed to polychlorinated biphenyls and dibenzofurans. *JAMA* 289:2943–2944.
- Karmaus W, DeKoning EP, Kruse H, Witten J, Osius N. 2001. Early childhood determinants of organochlorine concentrations in school-aged children. *Pediatr Res* 50:331–336.
- Kato Y, Haraguchi K, Shibahara T, Masuda Y, Kimura R. 1998. Reduction of thyroid hormone levels by methylsulfonyl metabolites of polychlorinated biphenyl congeners in rats. *Arch Toxicol* 72:541–544.
- Khan MA, Hansen LG. 2003. Ortho-substituted polychlorinated biphenyl (PCB) congeners (95 or 101) decrease pituitary response to thyrotropin releasing hormone. *Toxicol Lett* 144:173–182.
- Kimbro RD, Krouska CA. 2003. Human exposure to polychlorinated biphenyls and health effects: a critical synopsis. *Toxicol Rev* 22:217–233.
- Kimura-Kuroda J, Nagata I, Kuroda Y. 2005. Hydroxylated metabolites of polychlorinated biphenyls inhibit thyroid-hormone-dependent extension of cerebellar Purkinje cell dendrites. *Dev Brain Res* 154:259–263.
- Kleiman de Pisarev DL, Ferramola de Sancovich AM, Sancovich HA. 1995. Hepatic indices of thyroid status in rats treated with hexachlorobenzene. *J Endocrinol Invest* 18:271–276.
- Kleiman de Pisarev DL, Rios de Molina MC, San Martin de Viale LC. 1990. Thyroid function and thyroxine metabolism in hexachlorobenzene-induced porphyria. *Biochem Pharmacol* 39:817–825.
- Kleiman de Pisarev DL, Sancovich HA, Ferramola de Sancovich AM. 1989. Enhanced thyroxine metabolism in hexachlorobenzene-intoxicated rats. *J Endocrinol Invest* 12:767–772.
- Kohlmeier M. 1986. Direct enzymic measurement of glycerides in serum and in lipoprotein fractions. *Clin Chem* 32:63–66.
- Koopman-Esseboom C, Morse DC, Weisglas-Kuperus N, Lutke-Schipholt IJ, Van Der Pauw CG, Tuinstra LGMT, et al. 1994. Effects of dioxins and polychlorinated biphenyls on thyroid hormone status of pregnant women and their infants. *Pediatr Res* 36:468–473.
- Koppe JG, Bartonova A, Bolte G, Bistrup ML, Busby C, Butter M, et al. 2006. Exposure to multiple environmental agents and their effect. *Acta Paediatr Suppl* 95:106–113.
- Kramer MS, Chalmers B, Hodnett ED, Sevokskaya Z, Dzokovich I, Shapiro S, et al. 2001. Promotion of Breastfeeding Intervention Trial (PROBIT): a randomized trial in the Republic of Belarus. *JAMA* 285:413–420.
- Kuriyama S, Fidalgo-Neto A, Mathar W, Palavinskas R, Friedrich K, Chahoud I. 2003. Effect of low dose mono-ortho 2,3',4,4',5-pentachlorobiphenyl on thyroid hormone status and EROD activity in rat offspring: consequences for risk assessment. *Toxicology* 186:11–20.
- Kuriyama SN, Chahoud I. 2004. *In utero* exposure to low-dose 2,3',4,4',5-pentachlorobiphenyl (PCB 118) impairs male fertility and alters neurobehavior in rat offspring. *Toxicology* 202:185–197.
- Lackman GM, Schaller KH, Angerer J. 2004. Organochlorine compounds in breast-fed vs. bottle-fed infants: preliminary results at six weeks of age. *Sci Total Environ* 329:289–293.
- Lai TJ, Guo YL, Guo NW, Hsu CC. 2001. Effect of prenatal exposure to polychlorinated biphenyls on cognitive development in children: a longitudinal study in Taiwan. *Br J Psychiatry* 178:S49–S52.
- Langer P. 2005. Review: persistent organochlorinated pollutants (POPs) and human thyroid—2005. *Endocr Regul* 39:53–68.
- Langer P, Kocan A, Tajtakova M, Petrik J, Chovancova J, Drobná B, et al. 2005. Human thyroid in the population exposed to high environmental pollution by organochlorinated pollutants for several decades. *Endocr Regul* 39:13–20.
- Lans MC, Klasson-Wehler E, Willemsen M, Meussen E, Safe S, Brouwer A. 1993. Structure-dependent, competitive interaction of hydroxy-polychlorobiphenyls, -dibenzo-*p*-dioxins and -dibenzofurans with human transthyretin. *Chem Biol Interact* 88:7–21.
- Lans MC, Spiertz C, Brouwer A, Koeman JH. 1994. Different competition of thyroxine binding to transthyretin and thyroxine-binding globulin by hydroxy-PCBs, PCDDs and PCDFs. *Eur J Pharmacol* 270:129–136.
- Lanting CI, Fidler V, Huisman M, Boersma ER. 1998. Determinants of polychlorinated biphenyl levels in plasma from 42-month-old children. *Arch Environ Contam Toxicol* 35:135–139.
- Larsen PR, Kronenberg HM, Shlomo M, Polonsky KS. 2003. Williams Textbook of Endocrinology. Philadelphia, PA:W.B. Saunders Company.
- Li M-H, Hansen L. 1996a. Responses of prepubertal female rats to environmental PCBs with high and low dioxin equivalence. *Fundam Appl Toxicol* 33:282–293.
- Li M-H, Hansen LG. 1996b. Enzyme induction and acute endocrine effects in prepubertal female rats receiving environmental PCB/PCDF/PCDD mixtures. *Environ Health Perspect* 104:712–722.
- Liang QR, Liao RD, Su SH, Huang SH, Pan RH, Huang JL. 2003. Effects of lead on thyroid function of occupationally exposed workers [in Chinese]. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 21:111–113.
- Lopez CM, Pineiro AE, Nunez N, Avagnina AM, Villamil EC, Roses OE. 2000. Thyroid hormone changes in males exposed to lead in the Buenos Aires area (Argentina). *Pharmacol Res* 42:599–602.
- Mazhitova Z, Jensen S, Ritzén M, Zetterstrom R. 1998. Chlorinated contaminants, growth and thyroid function in schoolchildren from the Aral Sea region in Kazakhstan. *Acta Paediatr* 87:991–995.
- McNabb FM, Fox GA. 2003. Avian thyroid development in chemically contaminated environments: is there evidence of alterations in thyroid function and development? *Evol Dev* 5:76–82.
- Meeker JD, Altshul L, Hauser R. 2007. Serum PCBs, *p,p'*-DDE and HCB predict thyroid hormone levels in men. *Environ Res* 104:296–304.
- Miyazaki W, Iwasaki T, Takeshita A, Kuroda Y, Koibuchi N. 2004. Polychlorinated biphenyls suppress thyroid hormone receptor-mediated transcription through a novel mechanism. *J Biol Chem* 279:18195–18202.

- Morse DC, Groen D, Veerman M, Van Amerongen CJ, Koeter HBWM, Smits Van Prooije AE, et al. 1993. Interference of polychlorinated biphenyls in hepatic and brain thyroid hormone metabolism in fetal and neonatal rats. *Toxicol Appl Pharmacol* 122:27–33.
- Myers GL, Kimberly MM, Waymack PP, Smith SJ, Cooper GR, Sampson EJ. 2000. A reference method laboratory network for cholesterol: a model for standardization and improvement of clinical laboratory measurements. *Clin Chem* 46:1762–1772.
- Nagayama J, Okamura K, Iida T, Hiradawa H, Matsueda T, Tsuji H, et al. 1998. Postnatal exposure to chlorinated dioxins and related chemicals on thyroid hormone status in Japanese breast-fed infants. *Chemosphere* 37:1789–1793.
- National Academy of Clinical Biology. 2002. Laboratory Support for the Diagnosis and Monitoring of Thyroid Disease. Washington DC:National Academy of Clinical Biochemistry.
- National Cancer Institute. 1999. HHHQ-DietSys Analysis Software, version 4.02. Available: <http://appliedresearch.cancer.gov/DietSys/software.html> [accessed 7 May 2008].
- National Research Council. 2006. Fluoride in the Drinking Water: A Scientific Review of EPA's Standards. Washington DC:National Academies Press.
- Nawrot TS, Staessen JA, Den Hond EM, Koppen G, Schoeters G, Fagard R, et al. 2002. Host and environmental determinants of polychlorinated aromatic hydrocarbons in serum of adolescents. *Environ Health Perspect* 110:583–589.
- Negoita S, Swamp L, Kelley B, Carpenter DO. 2001. Chronic diseases surveillance of St. Regis Mohawk Health Service patients. *J Public Health Manag Pract* 7:84–91.
- Ness DK, Schantz SL, Moshtaghian J, Hansen LG. 1993. Effects of perinatal exposure to specific PCB congeners on thyroid hormone concentrations and thyroid histology in the rat. *Toxicol Lett* 68:311–323.
- Osius N, Karmaus W, Kruse H, Witten J. 1999. Exposure to polychlorinated biphenyls and levels of thyroid hormones in children. *Environ Health Perspect* 107:843–849.
- Patandin S, Dagnelie PC, Mulder PG, Op de Coul E, van der Veen JE, Weisglas-Kuperus N, et al. 1999. Dietary exposure to polychlorinated biphenyls and dioxins from infancy until adulthood: a comparison between breast-feeding, toddler, and long-term exposure. *Environ Health Perspect* 107:45–51.
- Persky V, Turyk M, Anderson HA, Hanrahan LP, Falk C, Steenport DN, et al. 2001. The effects of PCB exposure and fish consumption on endogenous hormones. *Environ Health Perspect* 109:1275–1283.
- Pluim HJ, de Vijlder JJ, Olie K, Kok JH, Vulsma T, van Tijn DA, et al. 1993. Effects of pre- and postnatal exposure to chlorinated dioxins and furans on human neonatal thyroid hormone concentrations. *Environ Health Perspect* 101:504–508.
- Pluim HJ, Koppe JG, Olie K, v d Slikke JW, Kok JH, Vulsma T, et al. 1992. Effects of dioxins on thyroid function in newborn babies. *Lancet* 339:1303.
- Porterfield SP. 1994. Vulnerability of the developing brain to thyroid abnormalities: environmental insults to the thyroid system. *Environ Health Perspect* 102:125–130.
- Porterfield SP, Hendrich CE. 1993. The role of thyroid hormones in prenatal and neonatal neurological development—current perspectives. *Endocr Rev* 14:94–106.
- Ribas-Fito N, Sala M, Cardo E, Mazon C, de Muga ME, Verdu A, et al. 2003. Organochlorine compounds and concentrations of thyroid stimulating hormone in newborns. *Occup Environ Med* 60:301–303.
- Rickenbacher U, McKinney JD, Oatley SJ, Blake CC. 1986. Structurally specific binding of halogenated biphenyls to thyroxine transport protein. *J Med Chem* 29:641–648.
- Robins JM, Cullen MR, Connors BB, Kayne RD. 1983. Depressed thyroid indexes associated with occupational exposure to inorganic lead. *Arch Intern Med* 143:220–224.
- Rozman K, Gorski JR, Rozman P, Parkinson A. 1986. Reduced serum thyroid hormone levels in hexachlorobenzene-induced porphyria. *Toxicol Lett* 30:71–78.
- Rylander L, Wallin E, Jonsson BA, Stridsberg M, Erfurth EM, Hagmar L. 2006. Associations between CB-153 and *p,p'*-DDE and hormone levels in serum in middle-aged and elderly men. *Chemosphere* 65:375–381.
- Sala M, Sunyer J, Herrero C, To-Figueras J, Grimalt J. 2001. Association between serum concentrations of hexachlorobenzene and polychlorobiphenyls with thyroid hormone and liver enzymes in a sample of the general population. *Occup Environ Med* 58:172–177.
- Sandau CD, Ayotte P, Dewailly É, Duffe J, Norstrom RJ. 2002. Pentachlorophenol and hydroxylated polychlorinated biphenyl metabolites in umbilical cord plasma of neonates from coastal populations in Québec. *Environ Health Perspect* 110:411–417.
- Sandstead HH. 1967. Effect of chronic lead intoxication on *in vivo* I-131 uptake by the rat thyroid. *Proc Soc Exp Biol Med* 124:18–20.
- Sandstead HH, Stant EG, Brill AB, Arias LI, Terry RT. 1969. Lead intoxication and the thyroid. *Arch Intern Med* 123:632–635.
- Sauer PJJ, Huisman M, Koopman-Esseboom C, Morse DC, Smits-van Prooije AE, van de Berg KJ, et al. 1994. Effects of polychlorinated biphenyls (PCBs) and dioxins on growth and development. *Hum Exp Toxicol* 13:900–906.
- Schell LM, DeCaprio A, Gallo MV, Hubicki L. 2002. Polychlorinated biphenyls and thyroid function in adolescents of the Mohawk Nation at Akwesasne. In: *Human Growth from Conception to Maturity* (Gilli G, Schell LM, Benso L, eds). London:Smith-Gordon, 289–296.
- Schell LM, Hubicki LA, DeCaprio AP, Gallo MV, Ravenscroft J, Tarbell A, et al. 2003. Organochlorines, lead, and mercury in Akwesasne Mohawk youth. *Environ Health Perspect* 111:954–961.
- Schumacher C, Brodtkin CA, Alexander B, Cullen M, Rainey PM, van Netten C, et al. 1998. Thyroid function in lead smelter workers: absence of subacute or cumulative effects with moderate lead burdens. *Int Arch Occup Environ Health* 71:453–458.
- Selkoff IJ, Hammond EC, Levin SM. 1983. Environmental Contaminants and the Health of the People of the St. Regis Reserve. Vol 1: Fluoride-Medical Survey Findings. New York:Environmental Sciences Laboratory, Mount Sinai School of Medicine of the City University of New York, 1–290.
- Seo BW, Li M-H, Hansen LG, Moore RW, Peterson RE, Schantz SL. 1995. Effects of gestational and lactational exposure to coplanar PCB congeners or TCDD on thyroid hormone concentrations in weanling rats. *Toxicol Lett* 78:253–262.
- Shrivastava VK, Katti SR, Sathyanesan AG. 1987. Effect of lead nitrate on thyroid function of the Indian palm squirrel, *Funambulus pennanti* (Wroughton). *Bull Environ Contam Toxicol* 38:981–984.
- Singh B, Dhawan D. 1999. Effect of lithium on thyroidal ¹³¹Iodine uptake, its clearance, and circulating levels of triiodothyronine and thyroxine in lead-treated rats. *Radiat Environ Biophys* 38:261–266.
- Singhal A, Cole TJ, Lucas A. 2001. Early nutrition in preterm infants and later blood pressure: two cohorts after randomised trials. *Lancet* 357:413–419.
- Sloan RJ, Jock K. 1990. Chemical Contaminants in Fish from the St. Lawrence River Drainage on Lands of the Mohawk Nation at Akwesasne and Near the General Motors Corporation/Central Foundry Division, Massena, NY Plant. Albany:New York State Department of Environmental Conservation.
- Smits JE, Fernie KJ, Bortolotti GR, Marchant TA. 2002. Thyroid hormone suppression and cell-mediated immunomodulation in American kestrels (*Falco sparverius*) exposed to PCBs. *Arch Environ Contam Toxicol* 43:338–344.
- Staessen JA, Nawrot T, Hond ED, Thijs L, Fagard R, Hoppenbrouwers K, et al. 2001. Renal function, cytogenetic measurements, and sexual development in adolescents in relation to environmental pollutants: a feasibility study of biomarkers. *Lancet* 357:1660–1669.
- Takser L, Mergler D, Baldwin M, de Grosbois S, Smargiassi A, Lafond J. 2005. Thyroid hormones in pregnancy in relation to environmental exposure to organochlorine compounds and mercury. *Environ Health Perspect* 113:1039–1045.
- Turyk ME, Anderson HA, Persky VW. 2007. Relationships of thyroid hormones with polychlorinated biphenyls, dioxins, furans, and DDE in adults. *Environ Health Perspect* 115:1197–1203.
- U.S. EPA. 1998. Guidance for Data Quality Assessment. Practical Methods for Data Analysis. Washington, DC:U.S. Environmental Protection Agency.
- Van Birgelen AP, Smit EA, Kampen IM, Groeneveld CN, Fase KM, van der Kolk J, et al. 1995. Subchronic effects of 2,3,7,8-TCDD or PCBs on thyroid hormone metabolism: use in risk assessment. *Eur J Pharmacol* 293:77–85.
- Van den BM, Birnbaum LS, Denison M, De VM, Farland W, Feeley M, et al. 2006. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci* 93:223–241.
- van Raaij JA, Kaptein E, Visser TJ, van den Berg KJ. 1993. Increased glucuronidation of thyroid hormone in hexachlorobenzene-treated rats. *Biochem Pharmacol* 45:627–631.
- van Raaij JA, van den Berg KJ, Engel R, Bragt PC, Notten WR. 1991a. Effects of hexachlorobenzene and its metabolites pentachlorophenol and tetrachlorohydroquinone on serum thyroid hormone levels in rats. *Toxicology* 67:107–116.
- van Raaij JA, van den Berg KJ, Notten WR. 1991b. Hexachlorobenzene and its metabolites pentachlorophenol and tetrachlorohydroquinone: interaction with thyroxine binding sites of rat thyroid hormone carriers *ex vivo* and *in vitro*. *Toxicol Lett* 59:101–107.
- Vreugdenhil HJ, Lanting CI, Mulder PGH, Boersma ER, Weisglas-Kuperus N. 2002. Effects of prenatal PCB and dioxin background exposure on cognitive and motor abilities in Dutch children at school age. *J Pediatr* 140:48–56.
- Wang S-L, Su PH, Jong S-B, Guo YL, Chou W-L, Papke O. 2005. *In utero* exposure to dioxins and polychlorinated biphenyls and its relations to thyroid function and growth hormone in newborns. *Environ Health Perspect* 113:1645–1650.
- WHO Collaborative Study Team on the Role of Breastfeeding on the Prevention of Infant Mortality. 2000. Effect of breastfeeding on infant and child mortality due to infectious diseases in less developed countries: a pooled analysis. *Lancet* 355:451–455.
- Wilson AC, Forsyth JS, Greene SA, Irvine L, Hau C, Howie PW. 1998. Relation of infant diet to childhood health: seven year follow up of cohort of children in Dundee infant feeding study. *Br Med J* 316:21–25.
- Zhou T, Ross DG, DeVito MJ, Crofton KM. 2001. Effects of short-term *in vivo* exposure to polybrominated diphenyl ethers on thyroid hormones and hepatic enzyme activities in weanling rats. *Toxicol Sci* 61:76–82.
- Zuurbier M, Leijts M, Schoeters G, ten Tusscher G, Koppe JG. 2006. Children's exposure to polybrominated diphenyl ethers. *Acta Paediatr Suppl* 95:65–70.